



UNIVERSITI PUTRA MALAYSIA

**ASSESSMENT OF FREE DIETARY GLUTAMATE INTAKE AMONG
MALAYS AND CHINESE IN THE KLANG VALLEY, MALAYSIA**

KHAIRUNNISAK BINTI MOHSIN

FSTM 2009 23

**ASSESSMENT OF FREE DIETARY GLUTAMATE
INTAKE AMONG MALAYS AND CHINESE IN
THE KLANG VALLEY, MALAYSIA**

KHAIRUNNISAK BINTI MOHSIN

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2009



**ASSESSMENT OF FREE DIETARY GLUTAMATE INTAKE AMONG
MALAYS AND CHINESE IN THE KLANG VALLEY, MALAYSIA**

By

KHAIRUNNISAK BINTI MOHSIN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science**

September 2009



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Master of Science

**ASSESSMENT OF FREE DIETARY GLUTAMATE INTAKE AMONG
MALAYS AND CHINESE IN THE KLANG VALLEY, MALAYSIA**

By

Khairunnisak Binti Mohsin

September 2009

Chairman : Azizah Abdul Hamid, PhD

Faculty : Food Science and Technology

Dietary intake assessment is a qualitative or quantitative evaluation of the degree of intake likely to happen, in which the estimation is made by multiplying the food consumption by the chemical concentration in the respective foods. In the present study, assessment of free dietary glutamate has been carried out among 360 Malay and Chinese adults in the urban area of Klang Valley, Malaysia. The food consumption data was obtained through a survey using food frequency questionnaires (FFQ) and condiments usage forms. In addition to that, the free glutamate content of food and condiments were analysed by High Performance Liquid Chromatography (HPLC) utilizing pre-column derivatization and fluorescence detection. Subsequently, using both data from survey and sample analysis, the study estimated free dietary glutamate intake of the respondents involved in the study. Results from the survey revealed that the average total food consumption by total respondents was 843 g/person/day, with significant ($p=0.017$) higher consumption among the Chinese (948 g/person/day) compared to the Malays (793 g/person/day). The study also showed that consumption of

condiments by both Malay and Chinese households was comparable, at 7.56 and 6.87 g/person/day, respectively. Chinese respondents, however, used more types of condiments compared to Malays. Meanwhile, results from sample analysis showed that the average free glutamate content in processed foods ranged from 0.34 ± 0.20 to 4.63 ± 0.41 mg/g. In food dishes, it was as low as 0.24 ± 0.15 mg/g in *roti canai* to 8.16 ± 1.99 mg/g in *dim sum*. Relatively, the free glutamate content was found to be higher in condiments at 0.28 ± 0 mg/g in mayonnaise to 786.00 ± 0 mg/g in monosodium glutamate. Calculated free glutamate intake from food by total respondents was 1.96 g/person/day; with intake by the Chinese was shown to be significantly ($p=0.001$) higher than Malays at 2.34 and 1.77 g/person/day, respectively. On the other hand, the average total free glutamate intake from condiments by total respondents was 0.27 g/person/day. The average total intake by the Malays, however, was significantly ($p=0.004$) higher than Chinese at 0.35 and 0.15 g/person/day, respectively. Overall, the average consumer need not be alarmed about the intake of glutamate from typical diet in their daily dietary pattern. Despite that, with heavy free glutamate intake (95th percentile) at more than 3.0 g/person/day may expose the consumers with possible adverse health effects.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENILAIAN PENGAMBILAN GLUTAMAT BEBAS MELALUI PEMAKANAN
DI KALANGAN KAUM MELAYU DAN CINA DI LEMBAH KLANG,
MALAYSIA**

Oleh

Khairunnisak Binti Mohsin

September 2009

Pengerusi : Azizah Abdul Hamid, PhD

Fakulti : Sains dan Teknologi Makanan

Penilaian pengambilan diet merupakan penilaian secara kualitatif atau kuantitatif terhadap tahap pendedahan pengambilan makanan. Anggaran ini dikira secara mendarabkan jumlah pengambilan makanan dengan kandungan bahan kimia yang terdapat di dalam makanan tersebut. Kajian penilaian pengambilan glutamat bebas di dalam makanan yang telah dijalankan ini melibatkan seramai 360 orang responden dewasa Melayu dan Cina di kawasan bandar, Lembah Klang, Malaysia. Data pengambilan makanan diperolehi melalui kaji selidik yang menggunakan borang soal selidik kekerapan pengambilan makanan (FFQ) dan penggunaan bahan perasa. Selain itu, kandungan glutamat bebas di dalam makanan dan bahan perasa telah dianalisis menggunakan alatan Kromatografi Cecair Berprestasi Tinggi (HPLC) berdasarkan terbitan pra-kolum dan pengesanan bersinar. Data daripada kedua-dua kaji selidik dan analisis sampel makanan kemudiannya digunakan untuk menganggar pengambilan

glutamat bebas di dalam makanan bagi responden yang terlibat di dalam kajian yang telah dijalankan. Keputusan daripada kaji selidik menunjukkan bahawa purata jumlah pengambilan makanan bagi kesemua responden adalah 843 g/orang/hari, dengan purata pengambilan makanan pada responden Cina (948 g/orang/hari) adalah ketara ($p=0.017$) lebih tinggi daripada responden Melayu (793 g/orang/hari). Kajian juga menunjukkan purata pengambilan bahan perasa bagi kedua-dua isi rumah Melayu dan Cina berada pada tahap yang sama, iaitu 7.65 dan 6.87 g/orang/hari masing-masing. Namun begitu, kaum Cina didapati lebih bervariasi dalam penggunaan bahan perasa berbanding kaum Melayu. Sementara itu, keputusan daripada analisis sampel makanan menunjukkan purata kandungan glutamat bebas di dalam makanan terproses berada diantara julat 0.34 ± 0.20 sehingga 4.63 ± 0.41 mg/g. Bagi hidangan makanan, julat kandungan glutamat bebas adalah 0.24 ± 0.15 mg/g dalam roti canai sehingga 8.16 ± 1.99 mg/g di dalam dim sum. Secara relatifnya, kandungan glutamat bebas adalah lebih tinggi dalam bahan perasa, iaitu 0.28 ± 0 mg/g di dalam mayonis sehingga 786.00 ± 0 mg/g di dalam monosodium glutamat. Anggaran pengambilan glutamat bebas daripada makanan bagi kesemua responden adalah 1.96 g/orang/hari; dengan pengambilan daripada kaum Cina adalah ketara ($p=0.001$) lebih tinggi berbanding kaum Melayu, iaitu pada 2.34 and 1.77 g/orang/hari masing-masing. Sementara itu, purata pengambilan glutamat bebas daripada bahan perasa bagi kesemua responden adalah sebanyak 0.27 g/orang/hari. Pengambilan di kalangan kaum Melayu adalah ketara ($p=0.004$) lebih tinggi daripada kaum Cina, iaitu sebanyak 0.35 dan 0.15 g/orang/hari masing-masing. Secara keseluruhannya, pengguna tidak perlu bimbang dengan pengambilan glutamat bebas daripada diet harian yang biasa. Walau bagaimanapun, pengambilan glutamat bebas

yang tinggi (pada 95 persentil) melebihi 3.0 g/orang/hari memungkinkan penggunaanya berhadapan dengan kesan sampingan kepada kesihatan.

ACKNOWLEDGEMENTS

Firstly, I thank Allah as this research has been successfully completed. Million thanks to my beloved parents and family for the unconditional love and supports all these while especially during difficult times throughout this study.

I also wish to extent my gratitude to my supervisor, Assoc. Prof Dr. Azizah Abdul Hamid for the guidance, invaluable advises and support; to my co-supervisor Prof Dr. Jinap Selamat for the opportunity and guidance; and special thanks goes out to Dr Nurul Izzah Ahmad for always guiding and lending me a helping hand during the hard times for the entire duration of the study.

Many thanks to IGTC and SEAAGS for funding the research, Dr. Akarat for monitoring the research activities and the staffs at Institute of Nutrition, Mahidol University, Thailand for the laboratory training assistance. Credits would also go out to all the laboratory assistants and science officer in Food Biochemistry laboratory and HPLC room.

Sincere appreciation extended to Saza, Farah and Redhuan for the help day and night during survey times, also in collecting and analysing the samples. To my lab mates in Food Safety and Quality laboratory especially Fatimah, Safzan, Hanis Izani, Kak Soffa, Kak Maimunah and Kak Afidah, thank you for the help, motivation and friendship. Last but not least, to all who have been involved directly or indirectly, I appreciate it much.

I certify that a Thesis Examination Committee has met on 15 September 2009 to conduct the final examination of Khairunnisak binti Mohsin on her thesis entitled “Assessment of Free Dietary Glutamate Intake among Malays and Chinese in the Klang Valley, Malaysia” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Mohhidin Othman, PhD

Senior Lecturer
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Roselina Karim, PhD

Senior Lecturer
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Internal Examiner)

Norimah Karim, PhD

Associate Professor
Faculty of Allied Health Sciences
Universiti Kebangsaan Malaysia
Malaysia
(External Examiner)

BUJANG BIN KIM HUAT, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:



This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science.

The members of the Supervisory Committee were as follows:

Azizah Abdul Hamid, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Jinap Selamat, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

Nurul Izzah Ahmad, PhD

Research Officer
Institute for Medical Research
(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 14 January 2010



DECLARATION

I hereby declare that the thesis is based on my original work except for quotation and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

KHAIRUNNISAK BINTI MOHSIN

Date:

TABLE OF CONTENTS

ABSTRACT	ii
ABSTRAK	iv
ACKNOWLEDGEMENTS	vii
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF APPENDICES	xvi
LIST OF ABBREVIATION	xviii

CHAPTER		
1	INTRODUCTION	1
2	LITERATURE REVIEW	
	2.1 Glutamic acid and glutamate	5
	2.2 Glutamate sources	6
	2.3 Monosodium glutamate (MSG)	7
	2.4 Applications of MSG and its usage levels in processed foods	8
	2.5 Trends of MSG consumption in various countries	11
	2.6 Safety evaluation of MSG	13
	2.7 MSG toxicology, sensitivity and adverse health effect	14
	2.8 Exposure (intake) assessment	16
	2.9 Exposure assessment approach: deterministic (point estimates) and probabilistic approach	17
	2.10 Dietary intake assessment to food additives	18
	2.11 Dietary intake assessment to glutamate	20
	2.11.1 Glutamate intake from food	20
	2.11.2 Glutamate intake from food additives	20
	2.12 Food consumption assessment	21
	2.12.1 Food supply data (food balance sheet)	24
	2.12.2 Household consumption surveys (budget survey)	24
	2.12.3 Individual dietary surveys	25
	2.12.4 Duplicate diets	27
	2.12.5 Biomarkers	28
	2.13 Food consumption survey in estimation of dietary exposure assessment	28
	2.13.1 Definition of FFQ	29
	2.13.2 Advantages and disadvantages of FFQ	30
	2.13.3 Development of FFQ	31

	2.13.4 Reproducibility of FFQ	32
	2.13.5 Validity of FFQ	33
2.14	Determination of free glutamate in food	
	2.14.1 Total diet study (TDS) method	35
	2.14.2 Analytical methods for determination of free glutamate	36
3	FOOD CONSUMPTION SURVEY AMONG MALAYS AND CHINESE IN KLANG VALLEY, MALAYSIA	
3.1	Introduction	39
3.2	Methodology	40
	3.2.1 General methodology for the study	40
	3.2.2 Food consumption survey	41
3.3	Results and Discussion	49
	3.3.1 Development and validation of the dietary tool	49
	3.3.2 Demographic profiles of the respondents involved in the main survey	52
	3.3.3 Average food consumption by Malay, Chinese and total respondents	55
	3.3.4 Average condiments consumption by the households	62
	3.3.5 Average condiments consumption by Malay and Chinese households	66
3.4	Conclusion	69
4	ANALYSIS OF FREE GLUTAMATE CONTENT IN PROCESSED FOOD, DISHES AND CONDIMENTS	
4.1	Introduction	71
4.2	Methodology	72
	4.2.1 Laboratory study	72
	4.2.2 Sampling procedure	72
	4.2.3 Location of the food and condiments sampling	75
	4.2.4 Sample collection and preparation	76
	4.2.5 Analysis method for determination of free glutamate in samples	76
	4.2.6 Statistical analysis	80
4.3	Results and discussion	81
	4.3.1 Recovery, LOD, LOQ and linearity test	81
	4.3.2 Free glutamate content of processed food and dishes	84
	4.3.3 Free glutamate content of condiments	89
4.4	Conclusion	92

5	ASSESSMENT OF FREE DIETARY GLUTAMATE INTAKE AMONG MALAYS AND CHINESE IN THE KLANG VALLEY, MALAYSIA	
5.1	Introduction	93
5.2	Methodology	95
5.2.1	Calculation for dietary intake assessment to free glutamate	95
5.2.2	Statistical analysis	95
5.3	Results and discussion	96
5.3.1	Free glutamate intake from food by Malay, Chinese and total respondents	96
5.3.2	Free glutamate intake from condiments by Malay, Chinese and total respondents’ households	100
5.4	Conclusion	104
6	CONCLUSION AND RECOMMENDATIONS	106
	REFERENCES	109
	APPENDICES	121
	BIODATA OF STUDENT	188
	LIST OF PUBLICATIONS	189

LIST OF TABLES

Table	Page
2.1 MSG usage levels in various processed foods	9
2.2 Recommended MSG usage levels	10
2.3 Daily intake of monosodium glutamate	12
2.4 Selected characteristics of the quantitative methods used in Europe to assess food consumption	22
3.1 Demographic information of respondents involved in FFQ survey (N=360)	53
3.2 Average consumption of each food groups (g/person/day) by Malay, Chinese and total respondents	56
3.3 Average condiments consumption by total respondents' households (N=302)	63
3.4 Average condiments consumption by Malay (N=188) and Chinese (N=114) households	67
4.1 Total food samples collected for free glutamate analysis	74
4.2 Recovery of added glutamic acid in food samples	82
4.3 Average free glutamate content in different type of foods consumed by Malay and Chinese respondents	85
4.4 Average free glutamate content of condiments	90
5.1 Average total free glutamate intake from food (mg/person/day) by Malay, Chinese and total respondents	97
5.2 Free glutamate intake from condiments by the households	101

LIST OF FIGURES

Figure		Page
2.1	Chemical structure of monosodium glutamate (MSG)	8
4.1	Chromatograms of HPLC analysis on free glutamate.	83

LIST OF APPENDICES

Appendix	Page
A1 Food frequency questionnaire (FFQ)	121
A2 Condiments Usage Form	129
A3a Questionnaire Tools- Food Photographs	134
A3b Questionnaire Tools- Condiments Photographs	144
A3c Questionnaire Tools- Standard measuring cups and spoons	150
A3d Questionnaire Tools- Digital food balance, sticker and basket to place the condiments	151
A4 Pictures during samples collections	152
B1 Respondent addresses as provided by Statistic Department of Malaysia	153
B2 Food items in pre-test FFQ	156
B3 Average food consumption in pilot study (N=30)	159
B4 Food items included in the finalized FFQ	162
B5a Chromatograms of HPLC analysis on free glutamate- condiment sample (shrimp paste (<i>belacan</i>))	164
B5b Chromatograms of HPLC analysis on free glutamate- food sample (rice noodle, fried (<i>mee-hoon goreng</i>))	165
B5c Chromatograms of HPLC analysis on free glutamate- food sample (chicken burger)	166
C1 Average consumption of each food items by total respondents (N=360)	167

C2	Average consumption of each food items by Malay (N=245) and Chinese (N=115) adults	172
C3	List of 50 highly consumed food items for analysis of free glutamate content	178
C4	List of condiments commonly used by the household and their dominant brand(s)	180
C5	Free glutamate intake by total respondents (N=360)	182
C6	Free glutamate intake by Malay (N=245) and Chinese (N=115) adults	185

ABBREVIATIONS

ADI	Acceptable daily intake
ANOVA	Analysis of variance
FAO	Food and Agricultural Organization
FBS	Food balance sheet
FFQ	Food frequency questionnaires
GEMS/Food	Global Environment Monitoring System/Food Contamination Monitoring and Assessment Programme
GRAS	Generally recognized as safe
JECFA	Joint FAO/WHO expert committee on food additives
LOAEL	Low observed adverse health effect
MANS	Malaysian adult nutrition survey
MOA	Ministry of Agriculture
MSG	Monosodium glutamate
NOAEL	No observed adverse health effect
SEAAGS	South East Asia Association of Glutamate Sciences
TDS	Total diet study
US FDA	United States Food and Drug Administration
WHO	World Health Organization

CHAPTER 1

INTRODUCTION

Glutamic acid is one of the most common, natural amino acid that occurs naturally in various foods either bound (part of protein) or in the free form. In foods it primarily exists in its salt form, glutamate (Federal register 1996). Glutamate is required for proper cell function and is a major excitatory transmitter within the brain (Watkins and Evans, 1981). Studies have shown that protein-rich foods like cheese, milk and meat consisted of substantial amount of bound-form glutamate, whereas vegetables like tomatoes and mushrooms contained high levels of free glutamate (Food Technologist's Expert Panel on Food Safety & Nutrition 1987). Neither the protein-bound glutamic acid nor the D-configuration has the same capability of the free L-glutamic acid, in terms of the flavour-enhancing activity. Consequently, only the free L-glutamic acid, through its salt monosodium glutamate, (MSG) has been used as a flavour enhancer.

MSG was first discovered in 1908, when Professor Kikunae Ikeda identified the unique taste attributed by glutamic acid as 'umami' or savoury, meat or broth-like taste (Ninomiya, 2001). MSG is the free form, sodium salt of glutamate. In its pure form it appears as a white, practically odorless, free-flowing crystalline powder; when dissolved in water (or saliva) it rapidly dissociates into free sodium and glutamate ions. When MSG is added to foods, it provides a similar flavoring function as the glutamate that occurs naturally in food. It is commonly used to improve the flavors of meats, poultry, seafood, snacks, soups and stews. It is also frequently been added to ready-to-

eat meals, flavorings, spices blends and vegetable. MSG is not only present in food as an added chemical, but also as a by-product of hydrolyzed vegetable proteins (HVP), which are widely used as seasonings and flavors in canned foods, dry mixes, sauces, and other manufactured products.

The U.S. Food and Drug Administration (FDA) has designated MSG as a Generally Recognised As Safe (GRAS) substance, along with many other common food ingredients such as salt, vinegar and baking powder. In addition, after many years of evaluations and meetings, the Joint FAO/WHO Expert Committee on Food Additives (JECFA) decided that MSG to be assigned as "Acceptable Daily Intake not specified" (JECFA, 1988). This indicates no toxicological concerns associated with its use as a food additive in accordance with good manufacturing practice (GMP). In 1991, the European Community's Scientific Committee for Food (SCF) confirmed the safety of MSG and came to the same conclusion that specification of an ADI was unnecessary (McKevith 2003). No "low observed adverse effect level" and "no observed adverse effect level" (LOAEL and NOAEL) has been set for added glutamate, at least by the US Committee on Dietary Reference Intakes (Panel on Micronutrient, 2002).

Despite these evaluations and findings, there has been considerable debate regarding the usage and safety issues of MSG for the past three decades. The source of these controversies started in 1968 when it was speculated that MSG could be the cause of adverse reactions following the ingestion of Chinese restaurant foods (Kwok, 1968). Since then, MSG has been implicated for eliciting a variety of symptoms, including

asthma (Allen *et al.* 1987; Hodge *et al.*, 1996), severe atopic dermatitis, (Van Bever *et al.*, 1989) headaches, facial burning, and abdominal pain (Yang *et al.*, 1997; Geha *et al.*, 2000). Hiroshi *et al.* (2002) reported that a diet with excess sodium glutamate over a period of several years may increase glutamate concentrations in the vitreous humour and may cause retinal cell destruction.

The issue arise in this case is whether high intakes might arise which could invalidate the assumption of safety at all foreseeable levels of exposure in food additive use. In addition to that, the assumption of absence hazard can be invalid with high intakes of MSG.

Different approach has been used to estimate the MSG intake around Southeast Asia countries; however, in many of the previous studies, the estimation of MSG consumption did not consider free glutamate that may exist in food. Therefore it is necessary that a complete study on the total exposure of the public to MSG and free glutamate from foods as well as from the condiments in Malaysia to be conducted. As it has not been comprehensively done, the population study using the exposure assessment approach has to be carried out.

Exposure assessment, as one of the crucial component of the risk assessment process, is defined as the qualitative and/or quantitative evaluation of the likely intake of biological, chemical or physical agents via food as well as exposure from other relevant sources (WHO, 1997). In the case of dietary exposure to glutamate it should be assessed

by combining data on food consumption with data on the glutamate concentration of the respective food. The term “dietary exposure” is used synonymously with the term “dietary intake” as both are acceptable and used concurrently (WHO, 2008).

In the present study, assessment to free dietary glutamate among the Malays and Chinese adults in Klang Valley, Malaysia is evaluated from food as well as from condiments. The output of the study will be a link between the safety information derived from animal experiment and the current measure to manage the risk that may be generated by glutamate consumption of general population including heavy users. It also could be used as the database for the next study conducted on the other part of Malaysia.

The objectives of the study are generally to determine the usage, consumption and potential exposure (intake) of consumers to MSG and free glutamate in food and food products. Specifically, the objectives are:

1. To determine the food consumption pattern, also the household use of MSG and glutamate-containing condiments.
2. To determine the free glutamate levels in ready-to-eat foods, processed foods, dishes and condiments available to consumers.
3. To evaluate the total exposure to glutamate from food consumption data and free glutamate content.

CHAPTER 2

LITERATURE REVIEW

2.1 Glutamic acid and glutamate

Glutamic acid was first identified from wheat gluten more than 100 years ago in Germany, but it was in Japan that its subtle effect on the taste of food was identified and given the name “umami” in 1908 by Professor Kikunae Ikeda. Ikeda was the first person who chemically isolated glutamate from dried kombu or sea tangle extract (*Laminaria Japanica*) (Ninomiya, 2001). He also suggested that free glutamate elicits a taste that is distinct from the four known primary tastes: sour, sweet, salty, and bitter. In China, the word *xianwei*, which represents the taste common in fish and meat, corresponds to umami. The same is true for savoury in English, *osmazome* in old French, and *gulih* in Indonesian (Matheis, 1999; Maga, 2002; Sugita, 2002; Prescott, 2004).

Glutamic acid, being one of the 20 amino acids that make up human proteins, has many important functions in human body. It is important for proper cell function, one of the building blocks in protein synthesis, a major starting material in the biosynthesis of other amino acids and also important in brain function as an excitatory neurotransmitter (Watkins and Evans, 1981). The average adult body contains approximately 10 g free glutamate (of which 2.3 g is in the brain) and 2 kg of the bound form. The turnover of